






How to recognise measles?



1. What is measles? Measles is an infectious disease that is caused by measles virus.



2. The source of infection: an ill man.




3. The routes of infection: airborne (runny nose, cough) and personal contact.

Measles symptoms:

High fever lasts for 7 days.

After a few days there is maculopapular rash on the face and neck.


At the beginning: runny nose, cough, conjunctivitis with photophobia, tiny spots in the inside of the cheeks and tongue.



In the next 3 days the rash changes its place down from the head to the torso and then to the hands and feet.

Prevention


- Compliance with the basic hygiene rules (washing and hand disinfection),
- The preventive vaccination (double vaccination immunizes fully; those who received only one dose of vaccination are recommended to vaccinate again).




The Provincial Sanitary—Epidemiological Station in Szczecin




How to recognise measles?



1. What is measles? Measles is an infectious disease that is caused by measles virus.



2. The source of infection: an ill man.




3. The routes of infection: airborne (runny nose, cough) and personal contact.

Measles symptoms:

High fever lasts for 7 days.

After a few days there is maculopapular rash on the face and neck.


At the beginning: runny nose, cough, conjunctivitis with photophobia, tiny spots in the inside of the cheeks and tongue.



In the next 3 days the rash changes its place down from the head to the torso and then to the hands and feet.

Prevention

- Compliance with the basic hygiene rules (washing and hand disinfection),
- The preventive vaccination (double vaccination immunizes fully; those who received only one dose of vaccination are recommended to vaccinate again).



The Provincial Sanitary—Epidemiological Station in Szczecin